




IL 04/000567

מדינת ישראל
STATE OF ISRAEL



Ministry of Justice
Patent Office

משרד המשפטים
לשכת הפטנטים

This is to certify that
annexed hereto is a true
copy of the documents as
originally deposited with
the patent application
particulars of which are
specified on the first page
of the annex.

זאת לתעודה כי
רצופים בזה העתקים
נכונים של המסמכים
שהופקדו לכתחילה
עם הבקשה לפטנט
לפי הפרטים הרשומים
בעמוד הראשון של
הנספח.

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

05-07-2004
This היום
רשם הפטנטים
Commissioner of Patents



נתאשר
Certified

BEST AVAILABLE COPY

לשימוש הלשכה
For Office Use

מספר : Number	156669
תאריך : Date	26-06-2003
הוקדם / נדחה Ante/Post-dated	

חוק הפטנטים, תשכ"ז - 1967
PATENT LAW, 5727-1967
ב ק ש ה ל פ ט נ ט
Application for Patent

אני, (שם המבקש, מענו ולגבי גוף מאגד - מקום התאגדותו)
I (Name and address of applicant, and in case of body corporate-place of incorporation)

Neurim Pharmaceuticals (1991) Ltd.
8 Hanechoshtet Street
Tel Aviv 69710
ISRAEL

נעורים פרמצבטיות (1991) בע"מ
רח' הנחושת 8
תל-אביב 69710

An Israeli Corporation

חברה ישראלית

ששמה הוא
Owner, by virtue of

הדין
the Law

בעל אמצאה מכח
of an invention the title of which is

נגזרי 2-אמינובנזויל

(בעברית)
(Hebrew)
(באנגלית)
(English)

2-AMINO BENZOYL DERIVATIVES

hereby apply for a patent to be granted to me in respect thereof.

מבקש בזאת כי ינתן לי עליה פטנט

בקשת חלוקה Application of Division	בקשת פטנט מוסף Application for Patent Addition	דרישה דין קדימה Priority Claim		
מבקשת פטנט from Application	לבקשה / לפטנט to Patent / Appl	מספר/סימן Number / Mark	תאריך Date	מדינת האגוד Convention Country
No. _____ מס' _____ dated _____ מיום _____	No. _____ מס' _____ dated _____ מיום _____			
יפוי כח: כללי / מיוחד / רצוף בזה / עוד יוגש P.O.A. : general / individual-attached / to be filed later הוגש בענין _____ 130169 filed in case				
המען למסירת מסמכים בישראל: ירמיהו מ. בן-דוד ושות' בע"מ ת.ד. 45087 ירושלים 91450 Address for Service in Israel: Jeremy M. Ben-David & Co. Ltd. POBox 45087 Jerusalem 91450				
חתימת המבקש Signature of Applicant		היום 26 בחודש יוני שנת 2003 This 26 day of June of the year 2003		

Our ref.: NEUR 202/13.1 מספרנו:

לשימוש הלשכה For Office Use

טופס זה, כשהוא מוטבע בחותם לשכת הפטנטים ומושלם במספר ובתאריך ההגשה, היינו אישור להגשת הבקשה שפרטיה רשומים לעיל.
This form, impressed with the Seal of the Patent Office and indicating the number and date of filing, certifies the filing of the application, the particulars of which are set out above.

נגזרי 2-אמינובנזויל

2-AMINO BENZOYL DERIVATIVES

Neurim Pharmaceuticals (1991) Ltd.

נעורים פרמצבטיות (1991) בע"מ

NEUR 202/13.1

2-Aminobenzoyl derivatives

נגזרי 2-אמינובנזויל

FIELD OF THE INVENTION

The present invention relates to novel 2-aminobenzoyl derivatives, pharmaceutical formulations containing them, and use of the compounds in the manufacture of medicaments for treating or preventing various diseases.

BACKGROUND OF THE INVENTION

The present invention relates generally to a novel compounds, their use in therapy, and pharmaceutical compositions containing them.

The amino acid tryptophan is converted biologically through the "kynurenine pathway" (Beadle et al. 1947, see Heidelberger et al. 1949). Over 95 % of all dietary tryptophan is metabolized to kynurenines (Wolf 1974). In peripheral tissues, in particular the liver, the indole ring of tryptophan is modified by either tryptophan dioxygenase or indoleamine 2,3-dioxygenase, which results in the formation of formylkynurenine. Kynurenine formylase then rapidly converts formylkynurenine to L-kynurenine, which is the key compound in the kynurenine pathway (Mehler & Knox 1950). L-kynurenine is present in low concentrations in the blood, the brain and in peripheral organs and it can easily cross the blood-brain barrier through the large neutral amino acid carrier. L-kynurenine is metabolized by three different enzymes in mammalian tissues: kynurenine 3-hydroxylase which form 3-hydroxy-kynurenine (3-HK); kynureninase which forms anthranilic acid and kynurenine aminotransferase (KAT) which causes the formation of kynurenic acid. 3-HK is metabolized by the same KAT to yield xanthurenic acid, a metabolically inert side product of the pathway, or by kynureninase to give rise to 3-hydroxyanthranilic acid, which is eventually converted to quinolinic acid. Finally, quinolinic acid is metabolized by quinolinic acid phosphoribosyltransferase, yielding nicotinic acid mononucleotide and subsequent degradation products including the end product NAD^+ .


Kynurenic acid, 3-hydroxykynurenine and quinolinic acid are all neuroactive intermediates of this catabolic cascade. 3-hydroxykynurenine is a free radical generator, which has been shown to cause induction of apoptosis, potentiation of excitotoxicity, cataract formation, neurodegenerative diseases, stroke, traumatic injury, neurovirological diseases and neuroinflammation. Quinolinic acid is an *N*-methyl-D-aspartate (NMDA) receptor agonist and free radical generator, and as such it can cause excitotoxicity, neurodegenerative diseases, stroke, traumatic brain injury, epilepsy, cerebral malaria, perinatal hypoxia, neurovirological diseases and neuroinflammation. Endogenous quinolinic acid might lead to NMDA receptor activation to promote excitotoxicity and neurotoxicity leading to physiological and pathological processes that are mediated by NMDA receptors. Among the three neuroactive kynurenines, kynurenic acid (KYNA) has recently received the most attention. First described as a neuroinhibitory compound two decades ago, KYNA, at high, nonphysiological, concentrations is a broad-spectrum antagonist of ionotropic glutamate receptors. High concentrations of KYNA are anticonvulsant and provide excellent protection against excitotoxic injury. At much lower concentrations, KYNA acts as a competitive blocker of the glycine coagonist site of the NMDA receptor and as a noncompetitive inhibitor of the $\alpha 7$ nicotinic acetylcholine receptor. The fact that the affinity of KYNA to these two Ca^{2+} -permeable receptors is in the range of KYNA levels in the human brain and reasonably close to the (lower) KYNA content of the rodent brain suggests a physiological function in glutamatergic and cholinergic neurotransmission. Direct support for such a role has been provided, for example, by *in vivo* studies in the rat striatum where a reduction in KYNA levels enhances vulnerability to an excitotoxic insult and, conversely, modest elevations of KYNA inhibit glutamate release (Schwarz and Pellicciari, 2002).

Since kynurenines have been suggested to participate not only in the pathophysiology of neurodegenerative and seizure disorders, but also to play a role in a large number of etiologically diverse CNS diseases, it is important to modulate their formation. We propose that the 2-aminobenzoyl derivatives (kynurenine-like compounds) described here will be useful for such therapeutic intervention. Suggested mechanism of action could be, but is not limited to, by

inhibiting enzymes in the kynurenine pathway, and/or inhibiting intermediate compounds, and/or inhibiting free radical formation.

SUMMARY OF THE INVENTION


The invention relates to compounds having the formula (I):



(I)

and stereoisomers and pharmaceutically acceptable salts thereof, wherein: A is C₁₋₆ alkylene; R, R₁ and R₂ are independently hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido, S-alkyl or alkylthiol; X is >C₁₋₆ alkylene, >C=O or >C=S or a single bond; and Y is hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido styryl which may be ring-substituted by up to four substituents independently selected from among hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, S-alkyl, alkylthiol or -COQ, where Q is hydroxy, C₁₋₆ alkoxy, amino, mono- C₁₋₆ alkylamino, di- C₁₋₆ alkylamino, hydroxylamino, C₁₋₄ alkoxyamino or aryl-C₁₋₄-alkoxyamino, but excluding (a) the compounds where simultaneously X is >C=O, Y is methyl, A is CH₂CH₂, R is 5-methoxy, R₁ is H or formyl and R₂ is H, and (b) the compounds where the moiety -A(R₂)-NH-X-Y is -CH₂CH(COQ)-NH₂ or -CH(haloalkyl)-CH(COQ)-NH₂.

Without prejudice to the generality of the compounds of the present invention, a sub-group of presently preferred compounds (formula II) is defined by the facts that R is hydrogen, methyl or methoxy, R₁ is hydrogen or formyl, R₂ is hydrogen or carboxyl, and R₃ is hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido, S-alkyl or alkylthiol, and stereoisomers and pharmaceutically acceptable salts thereof.



(II)

Another sub-group of the present compounds is defined by the facts that in the formula (I) X is 2-furyl, 2-dihydrofuryl, 2-tetrahydrofuryl or (2-R^o-COO-)phenyl, any of which may be substituted by 1-2 substituents selected from C₁₋₄ alkyl, C₁₋₄ alkoxy, OH, nitro, and Y is hydrogen or styryl which is ring-substituted by up to two substituents independently selected from among halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, OH, nitro, aryl, aryl-C₁₋₄ alkyl, or aryl-C₁₋₄ alkoxy, and stereoisomers and pharmaceutically acceptable salts thereof.

The present invention includes within its scope also the pharmaceutical compositions containing as an active substance therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof as well as any possible isomer, or mixture of isomers, covered by the formula (I) in association with one or more pharmaceutically acceptable diluents, preservatives, solubilizers, emulsifiers, adjuvants, excipients and carriers are those conventionally used in pharmaceutical and veterinary formulations. The

present pharmaceutical formulations may be adapted for administration to humans and/or animals.

The compounds of formula (I) are useful for treating and/or preventing, and/or minimizing neuronal loss associated with stroke, ischemia, CNS trauma, hypoglycemia and surgery, CNS disorders including neurodegenerative diseases such as Alzheimer's disease, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease and Down's syndrome, treating or preventing the adverse consequences of the overstimulation of the excitatory amino acids, psychiatric disorders, e.g., sleep disorders, epilepsy and other convulsive disorders, anxiety, psychiatric diseases, psychosis, senile dementia, multi-infarct dementia, chronic pain (analgesia), glaucoma, CMV retinitis, urinary incontinence, and inducing anesthesia, as well as enhancing cognition, and preventing opiate tolerance and withdrawal symptoms.

By way of further elaboration or explanation of conditions which it is presently contemplated may be amenable to treatment by administration of the present compounds, such conditions include impotence; cardiovascular disorders including hypertension, preventing blood coagulation, anti-inflammation, neuropathy, chronobiological-based disorders, e.g., jet lag, circadian sleep disorders such as delayed sleep syndrome, shift-work problems, and seasonal-related disorders e.g. seasonal affective disorder (SAD); endocrine indications, e.g., contraception and infertility, precocious puberty, premenstrual syndrome, hyperprolactinemia, and growth hormone deficiency; neoplastic diseases including e.g. cancer and other proliferative diseases (benign and tumor prostate growth), immune system disorders including AIDS, conditions associated with senescence, ophthalmological diseases, cluster headache, migraine, skin-protection, diabetes stabilization and weight gain disorders (leptin, obesity), and as an aid to animal breeding, e.g., regulation of fertility, puberty, pelage color.

DETAILED DESCRIPTION OF THE INVENTION

Without prejudice to the generality of the compounds of the present invention, in addition to sub-groups already mentioned above, another sub-group of presently preferred compounds is defined by the facts that in formula (I), X is

a 2,4-dinitrophenyl group, A is CH_2CH_2 or CH_2CHCOOH and R_2 and Y are each hydrogen.

Another sub-group of the present compounds is defined by the facts that in formula (I), R_2 is hydrogen and at least one of the following conditions applies, namely: R is 5-methoxy; and/or A is CH_2CH_2 or CH_2CHCOOH , and within these sub-groups, R_1 is preferably also hydrogen. Illustrative combinations of X and Y in compounds of the invention, particularly where $\text{R}_1 = \text{H}$, are the following:

X is $-\text{CO}-$ and Y is 2-furyl; or

X is $-\text{CO}-$ and Y is 2-tetrahydrofuryl; or

X is $-\text{CH}_2-$ and Y is 2-tetrahydrofuryl; or

X is $-\text{CO}-$ and Y is 2-acetoxyphenyl; or

X is $-\text{CO}-$ and Y is 3,4-dihydroxystyryl or 3,4-dihydroxycinnamoyloxy.

The present invention also includes within its scope pharmaceutical compositions containing as an active substance therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof as well as any possible isomer, or mixture of isomers, covered by the formula (I) in association with one or more pharmaceutically acceptable diluents, preservatives, solubilizers, emulsifiers, adjuvants, excipients and carriers are those conventionally used in pharmaceutical and veterinary formulations. The present pharmaceutical formulations may be adapted for administration to humans and/or animals.

The pharmaceutical formulation according to the invention is preferably characterized by at least one of the following features:

- (i) it is adapted for oral, rectal, parenteral, transbuccal, intrapulmonary (e.g. by inhalation) or transdermal administration;
- (ii) it is in unit dosage form, each unit dosage comprising an amount of said at least one member which lies within the range of 0.0025-1000 mg;
- (iii) it is a controlled release formulation, wherein said at least one member is released at a predetermined controlled rate.

For oral administration, the pharmaceutical formulations may be utilized as e.g. tablets, capsules, emulsions, solutions, syrups or suspensions. For parenteral administration, the formulations may be utilized as ampoules, or otherwise as suspensions, solutions or emulsions in aqueous or oily vehicles. The need for suspending, stabilizing and/or dispersing agents will of course take

account of the fact of the solubility or otherwise of the active compounds, in the vehicles that are used in particular embodiments. The formulations may additionally contain e.g. physiologically compatible preservatives and antioxidants. In the formulations for topical application, e.g. creams, lotions or pastes, the active ingredient may be mixed with conventional oleaginous or emulsifying excipients.

The pharmaceutical formulations may also be utilized as suppositories with conventional suppository bases such as cocoa butter or other glycerides. Alternatively, the formulations may be made available in a depot form which will release the active composition slowly in the body, over a pre-selected time period. Moreover, the compounds of the invention may be administered by using transbuccal, intrapulmonary or transdermal delivery systems.

Also combinations of the compounds of formula I as well as their salts with other active ingredients, especially other neuroleptics, thymoleptics, anxiolitics, tranquilizers, analgetics, antiparkinson's drugs (dopaminergic and non-dopaminergic drugs) or the like, fall within the scope of the present invention.


The compounds of the present invention are useful for treating and/or preventing, and/or minimizing neuronal loss associated with stroke, ischemia, CNS trauma, hypoglycemia and surgery, CNS disorders including neurodegenerative diseases such as Alzheimer's disease, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease and Down's syndrome, treating or preventing the adverse consequences of the overstimulation of the excitatory amino acids, psychiatric disorders, e.g., sleep disorders, epilepsy and other convulsive disorders, anxiety, psychiatric diseases, psychosis, senile dementia, multi-infarct dementia, chronic pain (analgesia), glaucoma, CMV retinitis, urinary incontinence, and inducing anesthesia, as well as enhancing cognition, and preventing opiate tolerance and withdrawal symptoms.

By way of further elaboration or explanation of conditions which it is presently contemplated may be amenable to treatment by administration of the present compounds, such conditions include impotence; cardiovascular disorders including hypertension, preventing blood coagulation, anti-inflammatory, neuropathy, chronobiological-based disorders, e.g., jet lag, circadian sleep disorders such as delayed sleep syndrome, shift-work problems,

and seasonal-related disorders e.g. seasonal affective disorder (SAD); endocrine indications, e.g., contraception and infertility, precocious puberty, premenstrual syndrome, hyperprolactinemia, and growth hormone deficiency; neoplastic diseases including e.g. cancer and other proliferative diseases (benign and tumor prostate growth); immune system disorders including AIDS; conditions associated with senescence; ophthalmological diseases; cluster headache, migraine; skin-protection, diabetes stabilization and weight gain disorders (leptin, obesity), and as an aid to animal breeding, e.g., regulation of fertility, puberty, pelage color.


The invention will be illustrated by the following Examples. The following examples are understood to be illustrative only and are not intended to limit the scope of the present invention in any way.

Example 1: 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine.



Kynuramine .2HBr (125 mg) is dissolved in 5 cc of absolute ethanol in a 50 cc flask. A solution of 2,4-dinitrofluorobenzene, 71 mg in EtOH 5 cc is then added. (a clear yellow solution is formed). After five minutes, 2 cc of a 10% NaHCO₃ solution are introduced drop wise in the flask. The reaction is left at room temperature overnight. The following morning the formed light yellow precipitate is filtered, washed with water and ethanol and dried in UHV obtaining 80 mg of product (approx. yield 63%).

Example 2: 3-(2-aminobenzoyl)-2-(2,4-dinitroanilino)propanoic acid.



L-Kynurenine (125 mg) is dissolved in 5 cc of absolute ethanol in a 50 cc flask. A solution of 2,4-dinitrofluorobenzene, 71 mg in EtOH 5 cc is then added. (a clear yellow solution is formed). After five minutes, 2 cc of a 10% NaHCO₃ solution are introduced drop wise in the flask. The reaction is left at room temperature overnight. The following morning the formed light yellow precipitate is filtered, washed with water and ethanol and dried in UHV obtaining 80 mg of product (approx. yield 71%).

The invention includes also pharmaceutically acceptable salts of the compounds of formula (I), as well as the possible isomers covered by the formula (I) both separately and in admixture.

BIOLOGICAL TESTING OF COMPOUNDS OF THE INVENTION

Experiment 1:

Evaluation of the anti-Parkinsonian activity using MPTP-treated mice with/without a subthreshold dose of L-Dopa

Animals: six month old male C57 BL/6 mice, weighing 22-25 g were used. Following arrival at the laboratory, the mice were allowed to acclimatise for two weeks in a room with controlled temperature (21±1°C), and a constant light-dark schedule (12 hr on/12 hr off, lights on between 06.00 and 18.00 hrs). Free access to food and water was maintained throughout. They were housed in groups of 12 animals and tested only during the hours of light (08.00-15.00 hrs). All testing was performed in a normally lighted room. Each test chamber (i.e.

activity test cage) was placed in a soundproofed wooden box with 12 cm thick walls and front panels and had a dimmed lighting.

Behavioral measurements and apparatus: An automated device, consisting of macrolon rodent test cages (40 x 25 x 15 cm) each placed within two series of infra-red beams (at two different heights, one low and one high, 2 and 8 cm, respectively, above the surface of the sawdust, 1 cm deep), was used to measure spontaneous and/or drug-induced motor activity of MPTP and control mice. The following parameters were measured: LOCOMOTION was measured by the low grid of infrared beams. Counts were registered only when the mouse in the horizontal plane, ambulating around the test-cage. REARING was registered throughout the time when at least one high level beam was interrupted, i.e. the number of counts registered was proportional to the amount of time spent rearing. TOTAL ACTIVITY was measured by a sensor (a pick-up similar to a gramophone needle, mounted on a lever with a counterweight) with which the test cage was constantly in contact. The sensor registered all types of vibration received from the test cage, such as those produced both by locomotion and rearing as well as shaking, tremors, scratching and grooming.

Behavioral measurements (locomotion, rearing and total activity): Twelve days after MPTP injections (2 x 40 mg/kg, s.c., 24 hr interval), the mice were injected i.p with 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine (0.3, 1, 3, 10 mg/kg) or vehicle (10% DMSO, 1% CMC) and immediately thereafter placed in the activity test chambers and their motor behaviour were monitored for 60 min. After 60 min, the mice were injected with 5mg/kg L-Dopa (s.c) and then replaced in the test chamber and activity measurements maintained for a further 240 min. Each dose was separated by two days, starting from the lowest dose.

Table 1 presents the mean (\pm standard deviations) locomotion, rearing and total activity counts of MPTP-treated and control mice administered either 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine or vehicle administered with a subthreshold dose of L-Dopa. 1% level of significance is represented by an asterisk (Tukey HSD test).

TREATMENT	LOCOMOTION	REARING	TOTAL ACTIVITY
Vehicle	1000±145	920±181	10937±2812
MPTP+vehicle	200±90	225± 72	4530± 937
MPTP+0.3mg/kg 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine	273±64	290± 73	5160±1093
MPTP+1mg/kg 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine	473±108*	582±145*	6250±625*
MPTP+3mg/kg 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine	510±107*	731±110*	6563±781*
MPTP+10mg/kg 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine	470±110*	619±102*	6250±625*

2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine had no effect at any dose the first 60-min period before L-Dopa, as compared to the MPTP-treated vehicle mice. However, 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine improved significantly all three behavioural parameters when administered together with a subthreshold dose of L-Dopa. 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine (1, 3 or 10 mg/kg) improved significantly the locomotion, rearing and total activity of MPTP-treated mice, as compared to the MPTP vehicle group. 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine administered to vehicle control animals had no effects in any variable.

Experiment 2:

Electrophysiological characterisation of NMDA-activated currents in freshly isolated hippocampal neurones of rat.

Isolation of hippocampal neurons: Wistar rats (12-14 days) were decapitated without anaesthesia and the hippocampus was removed. It was manually cut into slices (0.2-0.4mm), in a solution containing (mM): 150 NaCl; 5 KCl; 1.25 NaH_2PO_4 ; 2 CaCl_2 ; 2 MgCl_2 ; 26 NaHCO_3 ; 20 glucose. Slices were preincubated in this solution for 30 min at room temperature. The enzymatic treatment proceeded in the same solution with lower Ca^{2+} concentration (0.5mM) containing 0.4 mg/ml protease from *aspergillus oryzae*. The incubation in the enzyme solution proceeded at 32°C within 10 min. Slices were kept subsequently in enzyme-free solution containing normal Ca^{2+} concentration and used within 6-8h for obtaining isolated neurons. Throughout the entire procedure the solutions were continuously saturated with a 95% O_2 and 5% CO_2 gas mixture to maintain pH 7.4. For cell dissociation the slice was transferred into the extracellular solution containing (mM): 150 NaCl; 5 KCl; 2 CaCl_2 ; 10 N-2-hydroxyethylpiperazine-N'-2-ethanesulphonic acid (Hepes); pH adjusted with NaOH to 7.4. Single cells were isolated from CA and CA3 zones of hippocampal slices by vibrodissociation method. They had diameter 10-15 μm and preserved a small part of dendritic tree. After isolation they were usually suitable for recording for 1-2h.

Salines and chemicals: The contents of the extracellular solution was as follows (in mM): 130 NaCl, 5KCl, 2 CaCl_2 , 20 N-2-hydroxyethylpiperazine-N'-2-ethansulfonic acid (Hepes); 0.1 μM TTX, 10 μM glycine, 300 μM L-aspartate; pH was adjusted with NaOH to 7.4.

The contents of the intracellular solution were as follows (in mM): 110CSF, 20Tris-HCl (pH=7.2). L-aspartate and glycine solutions were prepared on the day of experiment.

The tested substance 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine was dissolved in DMSO.

Current recording and data analysis: The drug-containing solutions were applied by the fast "concentration clamp" method using "jumping table" set-up (Pharma Robot, Kiev). The currents were recorded with patch clamp technique in the

whole-cell configuration. Recording of the currents was performed using EPC-7 L/M patch-clamp amplifier.

NMDA-activated currents: The currents were filtered at 3kHz (three-pole active Bessel filter) digitally sampled at the rate 6000 μ s per point for NMDA activated currents. NMDA-induced transmembrane currents were measured in the presence of 10 μ M glycine and 300 μ M L-aspartate in the control and test solutions. The currents were recorded at holding potential -70 mV.

Calculations: The inhibition of current at different concentrations of the substance was averaged at least for 4 cells. The effect of substance was measured as the mean ratio I/I_0 where I was the current under the action of substance and I_0 was the current in control conditions. S.D. represents standard deviation.


The action of 10 μ M 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine on NMDA-activated currents:

2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine 10 μ M	INDIVIDUAL CELLS: I/I_0				\pm SD
	1	2	3	MEAN	
	92,77	77,15	80,17	83,36	

This experiment revealed that 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine has a blocking activity on NMDA receptors.

CLAIMS


1. A compound having the formula (I):



(I)

and stereoisomers and pharmaceutically acceptable salts thereof, wherein:
 A is C₁₋₆ alkylene; R, R₁ and R₂ are independently hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido, S-alkyl or alkylthiol; X is >C₁₋₆ alkylene, >C=O or >C=S or a single bond; and Y is hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido styryl which may be ring-substituted by up to four substituents independently selected from among hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, S-alkyl, alkylthiol or -COQ, where Q is hydroxy, C₁₋₆ alkoxy, amino, mono- C₁₋₆ alkylamino, di- C₁₋₆ alkylamino, hydroxylamino, C₁₋₄ alkoxyamino or aryl-C₁₋₄-alkoxyamino, but excluding (a) the compounds where simultaneously X is >C=O, Y is methyl, A is CH₂CH₂, R is 5-methoxy, R₁ is H or formyl and R₂ is H, and (b) the compounds where the moiety -A(R₂)-NH-X-Y is -CH₂CH(COQ)-NH₂ or -CH(haloalkyl)-CH(COQ)-NH₂.

2. A compound according to claim 1, having formula (II):



(II)

wherein R is hydrogen, methyl or methoxy, R_1 is hydrogen or formyl, R_2 is hydrogen or carboxyl, and R_3 is hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido, S-alkyl or alkylthiol, and stereoisomers and pharmaceutically acceptable salts thereof.

3. Compounds according to claim 1, where in formula (I), X is 2-furyl, 2-dihydrofuryl, 2-tetrahydrofuryl or (2- R^o -COO-)phenyl, any of which may be substituted by 1-2 substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, OH, nitro, and Y is hydrogen or styryl which is ring-substituted by up to two substituents independently selected from among halogen, C_{1-4} alkyl, C_{1-4} alkoxy, OH, nitro, aryl, aryl- C_{1-4} alkyl, or aryl- C_{1-4} alkoxy, and stereoisomers and pharmaceutically acceptable salts thereof.

4. Compounds according to claim 1, and stereoisomers and pharmaceutically acceptable salts thereof, where in formula (I), R_2 is hydrogen and at least one of the following conditions applies, namely:

R is 5-methoxy; and/or

A is CH_2CH_2 or $CH_2CHCOOH$; and/or

R_1 is hydrogen; and/or

X is a 2,4-dinitrophenyl group and Y is hydrogen.

5. Compounds according to claim 1, and stereoisomers and pharmaceutically acceptable salts thereof, where in formula (I), X and Y are selected in combination as follows:

X is -CO- and Y is 2-furyl; or

X is -CO- and Y is 2-tetrahydrofuryl; or

X is -CH₂- and Y is 2-tetrahydrofuryl; or

X is -CO- and Y is 2-acetoxyphenyl; or

X is -CO- and Y is 3,4-dihydroxystyryl or 3,4-dihydroxycinnamoyloxy.

6. Compounds according to claim 5, wherein at least one of the following conditions applies, namely:

R is 5-methoxy; and/or

A is CH₂CH₂ or CH₂CHCOOH; and/or

R₁ is hydrogen.

7. 3-(2-aminobenzoyl)-2-(2,4-dinitroanilino)propanoic acid, and stereoisomers and pharmaceutically acceptable salts thereof.

8. 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine, and pharmaceutically acceptable salts thereof.

9. A pharmaceutical composition containing a therapeutically effective amount of at least one compound as defined in claim 1, in association with at least one pharmaceutically acceptable ingredient selected from diluents, preservatives, solubilizers, emulsifiers, adjuvants, excipients and carriers.

10. A pharmaceutical formulation according to claim 9, which is further characterized by at least one of the following features:

(i) it is adapted for oral, rectal, parenteral, transbuccal, intrapulmonary or transdermal administration;


(ii) it is in unit dosage form, each unit dosage comprising an amount of said at least one compound which lies within the range of 0.0025-1000 mg;

(iii) it is a controlled release formulation, wherein said at least one compound is released at a predetermined controlled rate;

(iv) it comprises additionally at least one known therapeutically active ingredient selected from neuroleptics, thymoleptics, anxiolytics, tranquilizers, analgesics, and antiparkinson's drugs.

11. Use of at least one compound as defined in any one of claims 1-8, or a pharmaceutical formulation as defined in claim 9 or claim 10, for the manufacture of a medicament for treatment or prevention of a physiological condition selected from stroke, ischemia, CNS trauma, hypoglycemia and surgery, CNS disorders including neurodegenerative diseases overstimulation of the excitatory amino acids, psychiatric disorders, epilepsy and other convulsive disorders, anxiety, psychosis, senile dementia, multi-infarct dementia, chronic pain (analgesia), glaucoma, CMV retinitis, urinary incontinence, and for inducing anesthesia, enhancing cognition, and preventing opiate tolerance and withdrawal symptoms, impotence, cardiovascular disorders including hypertension, preventing blood coagulation, neuropathy, anti-inflammatory, chronobiological-related disorders, seasonal-related disorders, endocrine indications, contraception and infertility, precocious puberty, premenstrual syndrome, hyperprolactinemia, and growth hormone deficiency; neoplastic disease, other proliferative diseases (benign and tumor prostate growth), immune system disorders, conditions associated with senescence, ophthalmological diseases, cluster headache, migraine, skin-protection, diabetes stabilization and weight gain disorders, and for use in animal breeding.

For the Applicants,



Jeremy M. Ben-David & Co. Ltd.
NEUR 202/13.1

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.